arXiv:hep-ph/0501022v1 4 Jan 2005

Annihilation contribution and $B \to a_0\pi, f_0K$ decays.

D. Delepine* and J. L. Lucio M.†

Instituto de Física, Universidad de Guanajuato
Loma del Bosque # 103, Lomas del Campestre,
37150 León, Guanajuato; México

Carlos A. Ramírez‡

Depto. de Física, Universidad Industrial de Santander,
A.A. 678, Bucaramanga, Colombia

We analyze the decays $B^0 \to a_0^{\pm}\pi^{\mp}$ and $B^{-,0} \to f_0K^{-,0}$ and show that within the factorization approximation a phenomenological consistent picture can be obtained. We show that in this approach the O_6 operator provides the dominant contributions to the suppressed channel $B^0 \to a_0^+\pi^-$. When the a_0 is considered a two quark state, evaluation of the annihilation form factor using Perturbative QCD implies that this contribution is not negligible, and furthermore it can interfere constructively or destructively with other penguin contributions. As a consequence of this ambiguity, the positive identification of $B^0 \to \pi^+a_0^-$ can not distinguish between the two or four quark assignment of the a_0 [6]. According to our calculation, a best candidate to distinguish the nature of a_0 scalar is $Br(B^- \to \pi^0a_0^-)$ since the predictions for a four quark model is one order of magnitude smaller than for the two quark assignment. When the scalars are seen as two quarks states, simple theoretical assumptions based on $SU(2)$ isospin symmetry provide relations between different B decays involving one scalar and one pseudoscalar meson.

INTRODUCTION

B factories provides large samples of $B-\bar{B}$ mesons allowing the study of physical phenomena such as CP violation, the determination of the CKM mixing angles and the search for new physics[1, 2]. Clearly, hadronic physics will beneficiate of the high statistics achieved, and the study of processes with small branching ratios will be possible. The full understanding of the B physics is still lacking as well as a systematic first principles description of the phenomena involved. Instead different theoretical approaches are compared to data and assumptions such as factorization, or estimation of the relative size of different contributions (tree level, annihilation, penguins, final state interactions) can be tested. This can be achieved in processes where the dominant contributions are suppressed by symmetry or accidental cancellations.

The BABAR and Belle collaborations already reported precise measurements of non-leptonic B meson decays involving scalar mesons with branching ratio of order as low as 10^{-6}. Thus for example, for the $B^0 \to f_0K^0$ channel, besides the branching ratio the CP violating asymmetries are reported and, from the two pion spectrum, the authors are able to obtain the mass and width of the $f_0(980)$[3]. This is not the case for the $B \to a_0(980)\pi$ where branching fraction for given final states are reported -in particular $a_0^-\pi^+$- however in this case it is not possible to separate the B^0 from the $\bar{B}^0$ decays, unless a dominant decay mechanism is assumed[4]. In this context it is worth remarking that the $B \to a_0(980)\pi$ decay was suggested as a place where α, the weak mixing angle, could be measured through the CP violating asymmetries [5]. However, it was shown that the $B \to a_0^+\pi^-$ is suppressed by G parity and also by isospin, which implies that in the symmetry limit no CP violating asymmetry is expected to be experimentally accessible [6]. Thus, theoretical arguments support the idea that the $B^0 \to a_0^+\pi^-$ is strongly suppressed, so that

---

[*] Electronic address: delepine@fisica.ugto.mx
[†] Electronic address: lucio@fisica.ugto.mx
[‡] Electronic address: mpramir@netscape.net

the reported branching ratio can be identified with the dominant $B^0 \rightarrow a_0^- \pi^+$ decay. As a by product, in [6] the author concludes that the positive identification of $B^0/\bar{B}^0 \rightarrow a_0^{\pm}\pi^{\mp}$ is an evidence against the four quark assignment of a_0 or else, for the breakdown of perturbative QCD. Both, the suppression of the $B^0 \rightarrow a_0^+\pi^-$ channel as well as the potential evidence in favor or against a four quark state are rather interesting observation that deserves further analysis.

The low lying scalar sector of QCD represents a major challenge [8]. From the experimental point of view, the determination of the nature of existing states has not been achieved while from the theory side no consistent interpretation of the experimental data exists [9]. This is so even though a number of processes involving the appropriated final state in the kinematical region of interest have been analyzed. Thus, for example, $\phi \rightarrow \pi\pi\gamma, J/\Psi \rightarrow \phi\pi\pi, \phi KK$ and central production involve the $f_0(980)$ and $a_0(980)$ [10], whereas the di-pion in the $\Upsilon(nS) \rightarrow \Upsilon(mS)\pi\pi$ and $D \rightarrow \pi\pi\pi$ decays include the kinematical region where the $f_0(600)$ is expected to appear [11, 12, 13]. Unfortunately, although different processes are included in the analysis, data is not good enough to provide a clear picture of the scalars. In fact no consensus exist even on the fundamental intrinsic properties (mass and width) of the low lying scalar mesons [2, 12].

The appropriated theoretical description of non-leptonic B decays involving scalar mesons is important not only to understand the nature of the scalars but also because these must be considered as background to other processes of interest in B physics. Since scalars, vectors and tensors couple to two pseudoscalars, the following decays lead to the same final state: $B \rightarrow PV, B \rightarrow SP, B \rightarrow TP$ and $B \rightarrow PPP$, where S, V, T and P stand for scalar, vector, spin 2 and pseudoscalar meson respectively. B decays involving scalar mesons have been considered by a number of authors. Thus for example in [14] the tree level Hamiltonian and quark model calculation are used to predict branching ratios, while sum rules [15] and gluon-penguin dominance ($b \rightarrow sg$) are the basis to interpret the scalars produced in B decays in terms of glue balls [16], or still using QCD corrected Hamiltonian plus factorization [17] to propose evidence for two quark content of the $f_0(980)$.

In this note we analyze the $B^0 \rightarrow f_0(980)K$ and $B \rightarrow a_0(980)\pi$ decays using the factorization approximation. To this end we use the $\Delta B = 1$ weak Hamiltonian including QCD corrections to next to leading order. To evaluate the matrix elements we use values reported in the literature or model dependent estimates of the decay constants and form factors. In particular, the annihilation contribution is considered and evaluated using Perturbative QCD. This is important in estimating contributions previously neglected, and it is also relevant to quantify the statement in [6] regarding the four quark nature of the $a_0(980)$.

$\bar{B}^0 \rightarrow a_0^{\pm}\pi^{\mp}$ AND $B^{0,-} \rightarrow f_0 K^{0,-}$

Following the conventional approach [18, 19, 20, 21, 22], we start with the $\Delta B = 1$ effective Hamiltonian $H_{eff}(q = d, s)$

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \left[\lambda_{uq}(C_1 O_1^u + C_2 O_2^u) - \lambda_{tq} \left(\sum_{i=3}^{10} C_i O_i + C_g O_g \right) \right] + h.c. \tag{1}$$

where $\lambda_{q'q} = V_{q'b}V^*_{q'q}, q = d, s, q' = u, c, t, V_{ij}$ are the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. The Wilson coefficients, C_i, including next to leading order QCD corrections, are evaluated at the renormalization scale $\mu \simeq m_B/2$. We use the conventions and C_i values reported in [20]. It remains to evaluate the matrix elements between the states of interest.

$$A(B \rightarrow PS) = \; < PS|H_{eff}|B > \tag{2}$$

P and S stand for pseudoscalar and scalar meson respectively. In terms of the amplitude the branching ratio is given by:

$$Br(B \rightarrow PS) \simeq \tau_B \frac{G_F^2 |A(B \rightarrow PS)|^2}{32\pi m_B} \tag{3}$$

with τ_B the appropriated B meson lifetime ($\tau_{B^+} = 1.65 \cdot 10^{-12}s, \tau_{B^0} = 1.56 \cdot 10^{-12}s$).

The matrix elements are evaluated using the assumption of factorization. In that approximation the matrix elements of interest are given by:

$$\begin{aligned} A_{\bar{B}^0\to\pi^- a_0^+} \simeq\ & \lambda_{ud}(a_1 X^{\pi-}_{\bar{B}^0 a_0^+} + a_2 X^{\bar{B}^0}_{(a_0^+\pi^-)_u}) - \lambda_{td}\Bigg[\left(a_4 + a_{10} - \frac{(a_6+a_8)m_\pi^2}{\widehat{m}(m_b+m_u)}\right) X^{\pi^-}_{\bar{B}^0 a_0^+} \\ & + \left(2(a_3-a_5) + a_4 + \frac{a_9 - a_7 - a_{10}}{2} - \frac{(a_6 - a_8/2)m_B^2}{m_u(m_b+m_d)}\right) X^{\bar{B}^0}_{(a_0^+\pi^-)_u}\Bigg] \end{aligned} \tag{4}$$

$$\begin{aligned} A_{\bar{B}^0\to\pi^- a_0^+} \simeq\ & \lambda_{ud}(a_1 X^{a_0^-}_{\bar{B}^0\pi^+} + a_2 X^{\bar{B}^0}_{(a_0^-\pi^+)_u}) - \lambda_{td}\Big[(a_4+a_{10})X^{a_0^-}_{\bar{B}^0\pi^+} - 2(a_6+a_8)\tilde{X}^{a_0^-}_{\bar{B}^0\pi^+} \\ & + \left(2(a_3-a_5) + a_4 + \frac{a_9 - a_7 - a_{10}}{2} - \frac{(a_6 - a_8/2)m_B^2}{m_u(m_d+m_b)}\right) X^{\bar{B}^0}_{(a_0^-\pi^+)_u}\Big] \end{aligned} \tag{5}$$

$$\begin{aligned} A_{B^-\to\pi^0 a_0^-} \simeq\ & \lambda_{ud}(a_1(X^{a_0^-}_{B^-\pi^0} + X^{B^-}_{a_0^-\pi^0}) + a_2 X^{\pi_u^0}_{B^- a_0^-}) - \lambda_{td}\Big[(a_4+a_{10})X^{a_0^-}_{B^-\pi^0} \\ & -2(a_6+a_8)\tilde{X}^{a_0^-}_{B^-\pi^0} - \left(a_4 - \frac{3}{2}(a_9-a_7) - \frac{1}{2}a_{10} + \frac{(a_6+a_8)m_\pi^2}{m_u(m_b+m_d)}\right) X^{\pi_u^0}_{B^- a_0^-} \\ & + \left(a_4 + a_{10} + \frac{(a_6+a_8)m_B^2}{\widehat{m}(m_b+m_u)} X^{B^-}_{a_0^-\pi^0}\right)\Big] \end{aligned} \tag{6}$$

$$\begin{aligned} A_{B^-\to\pi^- a_0^0} \simeq\ & \lambda_{ud} a_1 (X^{\pi^-}_{B^- a_0^0} + X^{B^-}_{a_0^0\pi^-}) - \lambda_{td}\Bigg[(a_4 + a_{10} - \frac{(a_6+a_8)m_\pi^2}{\widehat{m}(m_b+m_u)})X^{\pi^-}_{B^- a_0^0} \\ & + \left(a_4 + a_{10} - \frac{(a_6+a_8)m_B^2}{\widehat{m}(m_b+\widehat{m})}\right) X^{B^-}_{a_0^0\pi^-} + (a_8 - 2a_6)\tilde{X}^{a_d^0}_{B^-\pi^-}\Bigg] \end{aligned} \tag{7}$$

$$\begin{aligned} A_{B^-\to f^0 K^-} \simeq\ & \lambda_{us} a_1 \left[X^{K^-}_{B^- f^0} + X^{B^-}_{f^0 K^-}\right] - \lambda_{ts}\Bigg[\left(a_4 + a_{10} - \frac{2(a_6+a_8)m_B^2}{(m_b+m_u)(m_s+m_u)}\right) X^{B^-}_{f^0 K^-} \\ & + \left(a_4 + a_{10} - \frac{2(a_6+a_8)m_K^2}{(m_u+m_s)(m_b+m_u)}\right) X^{K^-}_{B^- f^0} - (2a_6 + a_8)\tilde{X}^{f_s^0}_{B^- K^-}\Bigg] \end{aligned} \tag{8}$$

$$\begin{aligned} A_{\bar{B}^0\to f^0\bar{K}^0} \simeq\ & -\lambda_{ts}\Bigg[\left((a_4 - \frac{a_{10}}{2} - \frac{(2a_6 - a_8)m_B^2}{(m_b+m_d)(m_s+m_d)}\right) X^{\bar{B}^0}_{f^0\bar{K}^0} \\ & + \left(a_4 - \frac{a_{10}}{2} - \frac{(2a_6-a_8)m_K^2}{(m_s+m_d)(m_b+m_d)}\right) X^{\bar{K}^0}_{\bar{B}^0 f^0} - (2a_6 - a_8)\tilde{X}^{f_s^0}_{\bar{B}^0\bar{K}^0}\Bigg] \end{aligned} \tag{9}$$

where $\widehat{m} = (m_u + m_d)/2$. For future reference in Table 1 we quote the numerical values of the a_i coefficients. These expressions are obtained by inserting the vacuum between the currents in all possible ways, and a typical $X^a_{b,c}$ product of matrix elements is parameterized in terms of form factors and decay constant as:

$$X^{\bar{B}^o}_{\bar{K}^0 f_0} = \ <\bar{K}^0 f^0|(\bar{s}d)_L|0><0|(\bar{d}b)_L|\bar{B}^0> \ = f_B(m_{f^0}^2 - m_K^2)F_0^{f^0\bar{K}^0}(m_B^2) \tag{10}$$

In the appendix we define in detail all of the $X^a_{b,c}$. Let us start by summarizing our knowledge about the decay constants and form factors entering the calculation. We can classify these in four categories: Pseudoscalar decay constants (f_π, f_K, f_B). The values of the two former decay constants are taken from [4] while for the later we use f_B = 170 MeV[23]. The second kind are the scalar decay constants (f_a, f_f). For these we use published values estimated using theoretical arguments [7]. We then have form factors of the type $F_0^{SP}(m_B^2)$, evaluated at the m_B scale, *i.e.* calculable with perturbative methods. In Table (II), we quote the values we use. Finally we need the form factor $F_0^{BS}(m_P^2), F_0^{BP}(m_S^2)$ where S and P stand for a scalar or pseudoscalar meson. For the decay under

References	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9	a_{10}
[19]	1.039	0.084	40	-440	-120	-620	$0.7 - i$	$4.7 - 0.3i$	$-94 - i$	$-14 - 0.3i$
[18]	1.050	0.053	48	$-412 - 36i$	-45	$-548 - 36i$	$0.7 - i$	$4.7 - 0.3i$	$-94 - i$	$-14 - 0.3i$
[20]	1.046	0.024	$72 - 0.3i$	$-379 - 102i$	$-27 - 0.3i$	$-431 - 102i$	$-0.81 - 2.4i$	$3.3 - 0.8i$	$-92.4 - 2.41i$	$0.34 - 0.8i$
[21]	1.061	0.011	63	-317	-60	-473	4	5.4	-87	-2.4
[22]	1.029	0.103	36	-228	-24	-298	12	7.6	-82	-8.2

TABLE I: Numerical values for the effective coefficientes a_i^{eff} for $b \to d$ transitions at scale $\mu \approx m_b$ (for $a_3, ..., a_{10}$ in units of 10^{-4} $(a_{2i-1} = C_{2i-1} + C_{2i}/N,\ a_{2i} = C_{2i} + C_{2i-1}/N)$)

f_B	170 MeV	[23]
f_K	159.8 MeV	[2]
f_π	130.7 MeV	[2]
$\tilde{f}_{f_s^0}$	180 MeV	[7]
$\tilde{f}_{a_0}$	400 MeV	[7]
$F_0^{B^0\pi^-}$	0.28	[24]
$F_0^{B^0K^-}$	0.34	[24]

TABLE II: Numerical values of the form factors

consideration these are evaluated at most at the scalar scale (around 1 GeV), *i.e.* these can not be computed using perturbative methods and, in general, few is known about their values.

For the $B \to f_0 K^0, B^- \to f_0 K^-$ and $B \to a_0 \pi$ we require $F_0^{B\pi}$, F_0^{BK}, $F_0^{Bf^0}$, $F_0^{Ba_0}$, $F_0^{a\pi}$ and $F_0^{f^0K}$. The two first ($F_0^{B\pi}$, F_0^{BK}) are relatively well-known and we shall use the value presented in refs.[24]. In principle, the other four form factors could be determined using experimental data[1] [3, 4, 5] given in Table (III) but since the experimental data involve large error bars, we prefer to evaluate some of these form factors. In this paper we focus our interest on the annihilation effects, for that reason we present an estimate of $F_0^{a_0\pi}$ using perturbative QCD and considering a_0 as a two or four quark state.

ANNIHILATION FORM FACTORS FROM PERTURBATIVE QCD.

At tree level the $\bar{B}^0 \to \pi^+ a_0^-$ is strongly suppressed due to the absence of second class currents. To get an evaluation for such a decay an estimate of the contribution of the B annihilation is necessary. The annihilation amplitude is proportional to $X^{\bar{B}^0}_{(a_0^-\pi^+)_u}$ which itself is proportional to the $F_0^{a_0^-\pi^+}(m_B^2)$ form factor. At the scale m_B^2, perturbative QCD provides an adequate framework to evaluate this form factor. So, below we compute this form factor using the standard approach of perturbative QCD [25] assuming the scalar meson a_0^- is a two quark state.

In order to fix the convention, we recall the form factor definitions :

$Br(\bar{B}^0 \to \pi^\pm a_0^\mp)$	$(2.8^{+1.65}_{1.47})\ 10^{-6}$	[4]
$Br(\bar{B}^0 - \to \pi^- a_0^0)$	$(3.6^{+2.25}_{-2.06}\ 10^{-6}$	[4]
$Br(B^- \to K^- f^0)$	$(13.5^{+3.6}_{-4.2})\ 10^{-6}$	[3]
$Br(\bar{B}^0 \to K^0 f^0)$	$(8.8 \pm 2.38)\ 10^{-6}$	[3]

TABLE III: Branching ratios of measured PS channel decays of B mesons

---

[1] we used $Br(f^0 \to 2\pi) = 0.68$ [2] in order to get the $Br(B^{-,0} \to K^{-,0} f^0)$ from published results.

$$\langle M_2(p_2)|L^u|M_1(p_1)\rangle = \left(p_1+p_2-\frac{m_1^2-m_2^2}{q^2}\right)_u F_+^{M_2M_1}(q^2)+\left(\frac{m_1^2-m_2^2}{q^2}\right)q_u F_0^{M_1M_2}(q^2), \tag{11}$$

$$\tag{12}$$

with $q=p_1-p_2$. Projecting the amplitude on q one obtains:

$$q_u\langle M_2(p_2)|L^u|M_1(p_1)\rangle = (m_1^2-m_2^2)F_0^{M_1,M_2}(q^2) \tag{13}$$

$$q_u\langle M_2(p_2)M_1(p_1)|L^u|0\rangle = (m_2^2-m_1^2)F_0^{M_2,M_1}(q^2) \tag{14}$$

$PQCD$ contributions to both amplitudes have exactly the same structure, so we compute $q_u\langle M_2(p_2)|L^u|M_1(p_1)\rangle$ following ref. [26] then $q_u\langle M_2(p_2)|L^u|M_1(p_1)\rangle$ is obtained just changing the sign of p_2. The form factors are expressed in terms of the distribution amplitudes:

$$\Psi_\pi(x,p) = \frac{-iI_c}{\sqrt{2N_c}}\phi_\pi(x)(\hat{p}+m_\pi)\gamma_5 \tag{15}$$

$$\Psi_{a_0}(x,p) = \frac{I_c}{\sqrt{2N_c}}\phi_{a_0}(x)(\hat{p}+m_{a_0})\gamma_5 \tag{16}$$

where I_C is the identity in color space, $\hat{p}=\ \gamma_\mu p^\mu$ and

$$\int \phi_\pi(x)dx = \frac{1}{2\sqrt{2N_c}}f_\pi \tag{17}$$

$$\int \phi_{a_0}(x)dx = \frac{1}{2\sqrt{2N_c}}f_{a_0} \tag{18}$$

The wave functions $\phi_{\pi,a_0}(x)$ are given by [27].

$$\phi_\pi(x) = \frac{2N_c}{2\sqrt{2N_c}}f_\pi x(1-x)+\cdots \tag{19}$$

$$\phi_{a_0}(x) = \frac{2N_c}{2\sqrt{2N_c}}f_{a_0}x(1-x)\left(1+B_1C_1^{3/2}(2x-1)\right)+\cdots \tag{20}$$

where $f_\pi=130MeV, f_{a_0}=1MeV, |B_1f_{a_0}|\simeq 75MeV$, and $C_1^{3/2}(2x-1)$ is the Gegenbauer polynomial. Thus, the matrix element is expressed as:

$$\begin{aligned} q_\mu\langle\pi(p_2)|L^\mu|a_0(p_1)\rangle &= -C(R)\frac{T_r(I_C)}{2N_c}g_s^2\int dxdy\phi_{a_0}(x)\phi_\pi(y) \\ &\left\{\frac{Tr\left[\gamma_5(\hat{p}_2+m_\pi)\gamma_\nu\hat{P_{1l}}q_\mu L^\mu(\hat{p}_1+m_{a_0})\gamma^\nu\right]}{k^2P_{1l}^2}+\right. \\ &\left.\frac{Tr\left[\gamma_5(\hat{p}_2+m_\pi)q_\mu L^\mu\hat{P_{2l}}\gamma^\nu(\hat{p}_1+m_{a_0})\gamma_\nu\right]}{k^2P_{2l}^2}\right\} \end{aligned} \tag{21}$$

where $C(R)=4/3, (p_1-p_2)^2=q^2=m_B^2$, $k=-xp_1+(1-y)p_2$, $P_{1l}=k+yp_2$, $P_{2l}=-k+(1-x)p_1$ and

$$P_{1l}^2 = x^2m_{a_0}^2+m_\pi^2+x(m_B^2-m_{a_0}^2-m_\pi^2), \tag{22}$$

$$P_{2l}^2 = (1-y)m_B^2+ym_{a_0}^2-m_\pi^2y(1-y). \tag{23}$$



FIG. 1: Allowed region for $F_0^{B^0 a^+}$ and $F_0^{\pi a}$ at one σ using experimental data on $Br(B^0 \to \pi^- a_0^+)$ and $Br(B^- \to \pi^- a_0^0)$

Integrating numerically, one gets

$$\left| F_0^{a_0^- \pi^+}(m_B^2) \right| \approx 0.004 \tag{24}$$

It is important to notice that the CP-conserving phase of the annihilation contributions is not fixed since we only known the absolute value of B_1.

GENERAL FRAMEWORK TO PREDICT $B \to PS$

One can proceed along similar lines for processes involving a_0 or f_0 scalar mesons. Instead, we use isospin, $SU(2)$ quark symmetry and the quark contents of the scalar mesons to obtain relation between the form factors. We also used available experimental data to obtain constraints on the form factor values. It turns out that the consistency of the two sets of values so obtained provide further confidence on the approach.

We assume the conventional quark content of the pseudo scalar mesons[2] and parameterize the mixing in the scalar sector in the strange-nonstrange basis as:

$$\sigma = \cos\phi_S \bar{n}n - \sin\phi_S \bar{s}s, \tag{25}$$

$$f_0 = \sin\phi_S \bar{n}n + \cos\phi_S \bar{s}s, \tag{26}$$

where $\bar{n}n = (\bar{u}u + \bar{d}d)/\sqrt{2}$, and the singlet-octet mixing angle θ_S is related to ϕ_S by $\phi_S - \theta_S = \cos^{-1}[1/\sqrt{3}] \simeq 55°$. A diagrammatic analysis of the contributions of the form factors, which involve the quark composition and isospin and $SU(2)$ symmetry between up and down quarks, lead the following relations :

$$\frac{\sqrt{2}}{\cos\phi_S} F_0^{B^- \sigma} = \frac{\sqrt{2}}{\sin\phi_S} F_0^{B^- f_0} = \frac{\sqrt{2}}{\sin\phi_S} F_0^{\bar{B}^0 f_0} = F_0^{\bar{B}^0 a_0^+} \tag{27}$$

From these relations it follows that $|F_0^{\bar{B}^0 f_0}| < |F_0^{\bar{B}^0 a_0^+}|/\sqrt{2}$.



FIG. 2: Allowed region for parameters $F_0^{B_0 f_0}$ and $F_0^{K^0 f_0}$ at one σ

It is possible to obtain similar relations between the annihilation form factors ($F_0^{a_0\pi}$ and $F_0^{f^0K}$), however we will not use $SU(3)$ symmetry since large deviations from the symmetry limit are expected. Using the experimental results given in Table (III), the values for the scalar masses given in ref.[2] and the values given in table (II) for the other form factors appearing in the amplitudes, it is possible to determine separately for each scalars a_0 and f_0 the allowed regions for the values of the form factors $F_0^{a_0\pi}$ ($F_0^{f_0K}$) and $F_0^{Ba_0}$ (F_0^{BK}) respectively.

The results are summarized in figures 1 and 2. Assuming that Perturbative QCD leads us the right order of magnitude for $F_0^{a_0\pi}$, it follows that:

$$0.14 \leq |F_0^{Ba_0}| \leq 0.21 \tag{28}$$

We should note that this result is compatible (even if slightly smaller) with the predictions for $F_0^{\bar{B}^0 a_0^+}(0) = 0.55 \pm 0.22$ [28] obtained in a model-dependent way. Using Eq.(27), it follows that

$$|F_0^{\bar{B}^0 f_0}| \leq 0.20 \tag{29}$$

Figure (2) shows the values allowed by the experimental data when one standard deviation is considered. One observes that $|F_0^{\bar{B}^0 f_0}| \leq 0.20$ requires a large contribution from $|F_0^{f_0K}|$. In fact, the smallest value for $|F_0^{f_0K}|$is around 0.05, which is more than one order of magnitude bigger than the PQCD evaluation of $|F_0^{a_0\pi}|$. It is interesting to note in this respect that:

$$\frac{|F_0^{f_0K}|}{|F_0^{a_0\pi}|} \approx \frac{m_K^2}{m_\pi^2} \approx 12 \tag{30}$$

Scalar mesons as $qq\bar{q}q$ states.

Several models where the scalars are four quark states [13, 29, 30, 31] have been published but no model is favored at present time. We shall apply our method to one example, following [29] we assume that the quarks contents of the scalars is given by



FIG. 3: Branching ratio for $Br(\bar{B}^0 \to \pi^+ a_0^-)$. The dot-dashed line correspond to the four quark assignement for a_0, and the space between both solid horizontal line is the value for $Br(\bar{B}^0 \to \pi^+ a_0^-)$ expected from 2 quark models for a_0

$$\begin{aligned} a_0^+ &= uu\bar{d}\bar{s}, \quad a_0^- = ds\bar{u}\bar{s}, \quad a_0^0 = \frac{1}{\sqrt{2}}\left(us\bar{u}\bar{s} - ds\bar{d}\bar{s}\right), \quad K_0^+ = ud\bar{d}\bar{s}, \quad K_0^0 = ud\bar{u}\bar{s}, \quad \bar{K}_0^0 = us\bar{u}\bar{d}, \quad K_0^- = ds\bar{u}\bar{d} \\ f_0 &= \frac{\cos\phi}{\sqrt{2}}\left(su\bar{s}\bar{u} + sd\bar{s}\bar{d}\right) + \sin\phi\ ud\bar{u}\bar{d}, \quad \sigma = -\frac{\sin\phi}{\sqrt{2}}\left(su\bar{s}\bar{u} + sd\bar{s}\bar{d}\right) + \cos\phi\ ud\bar{u}\bar{d} \end{aligned} \tag{31}$$

where the mixing angle is obtained from the relation $\tan\phi = -0.19$ (for $m_\sigma = 0.47$ GeV), so $\phi = -5.4^{\mathrm{o}}$ and 84.6^{o}.

It is well-known that perturbative QCD predicts that the form factor will go like $1/q^{2(n-1)}$ where n is the number of constituents of the hadron. If $n = 4$, $F_0^{a_0^-\pi^+}(m_B^2)$ is strongly suppressed and annihilation can be neglected. Varying the experimental results within one σ and using the fact that in four quark models for scalars the annihilation does not contribute to the processes ($F_0^{a_0\pi} = F_0^{Kf_0} = 0$), one concludes that:

$$0.70 \leq F_0^{Bf_0} \leq 0.75 \tag{32}$$

Proceeding in the same way with processes $B \to a_0^0\pi^-, a_0^\pm\pi^\mp$, one gets

$$0.15 \leq |F_0^{Ba_0}| \leq 0.20 \tag{33}$$

which are closed to the values of $|F_0^{Ba_0}|$ obtained assuming the scalars are two quark states.

Sub-dominant processes $Br(\bar{B}^0 \to \pi^+ a_0^-)$ and annihilation

Once the allowed values for the form factors $F_0^{Bf_0}$ and $F_0^{Ba_0}$ have been constrained, we turn our attention to the subdominant processes $Br(\bar{B}^0 \to \pi^+ a_0^-)$ which is strongly suppressed by G parity and isospin. In [6] the author concludes that a positive identification of this process is an evidence against the four-quark assignment of a_0 or else for breakdown of perturbative QCD. Using our estimates for annihilation contributions obtained using PQCD one can predict the values of $Br(\bar{B}^0 \to \pi^+ a_0^-)$. The results are presented in figure 3. In the four quark models for a_0 where annihilation is strongly suppressed, one gets for $Br(\bar{B}^0 \to \pi^+ a_0^-) \approx 10^{-7}$. In the two quark model for a_0, the main source of uncertainty is the phase of the annihilation contributions which cannot be fixed using PQCD. Varying between 0 and π the CP-conserving phase for the annihilation, it follows that:

$$10^{-9} \leq Br(\bar{B}^0 \to \pi^+ a_0^-) \leq 4 \times 10^{-7} \tag{34}$$

where the lower limit is obtained when annihilation interferes destructively and the upper limit when annihilation interferes constructively with the other contributions. Our conclusion is that this B decay cannot be used to distinguish between two and four quark assignment of the a_0, unless one can obtain an independent determination of the annihilation phase.

Another channel suppressed by G parity is $B^- \to \pi^0 a_0^-$. It is interesting that this channel is better to distinguish between the 2 or 4 quark models for a_0. Indeed, in 4 quark models for a_0, with the value for $|F_0^{Ba_0}|$ determined in previous sections, one gets

$$2 \times 10^{-9} \leq Br(B^- \to \pi^0 a_0^-) \leq 10^{-8} \tag{35}$$

while in the 2 quark model:

$$6.4 \times 10^{-8} \leq Br(B^- \to \pi^0 a_0^-) \leq 2.4 \times 10^{-7} \tag{36}$$

CONCLUSIONS

In this paper we consider processes for which the dominant contribution is suppressed. Using the factorization approximation and available experimental data, we estimate the effect of annihilation penguins contribution to the processes $Br(\bar{B}^0 \to \pi^\pm a_0^\mp)$ and $Br(\bar{B}^{0,-} \to K^{0,-} f_0)$. We show that a consistent picture can be obtained when the scalars are described as two quark states, although one requires an important contributions from annihilation penguins to $Br(\bar{B}^{0,-} \to K^{0,-} f_0)$. Within our analysis the four quark models for f_0 cannot be excluded.

Applying our estimates of the annihilation contributions to suppressed processes like $Br(B^- \to \pi^0 a_0^-)$ and $Br(\bar{B}^0 \to \pi^+ a_0^-)$, we conclude that the positive identification of $\bar{B}^0 \to \pi^+ a_0^-$ cannot be taken as evidence for the four quark assignment of a_0. This is in contrast with ref.[6], where the annihilation contribution is not quantified. Relevant for this conclusion is the ambiguity in the CP-conserving phase of the annihilation penguins contributions. Our best candidate process to distinguish the nature of a_0 scalar is $Br(B^- \to \pi^0 a_0^-)$ where the predictions for 4 quark models are typically one order of magnitude smaller than 2 quark models.

Using the mesons quark content, $SU(2)$ quark symmetry and isospin we derive relations between the form factors $F_0^{Bf_0}$ to $F_0^{Ba_0}$ for different charge states. One can extend the analysis to $SU(3)$ however one expects large deviations from the symmetry limit. This restricts the applicability of our approach to the four quark states since in that kind of models scalars necessarily involve strange quarks.

---

[1] P. Ball et al., hep-ph/0003238, http://lhcb.web.cern.ch/lhcb; K. Anikeev et al., hep-ph/0201071, http://www-btev.fnal.gov; http//www.hera-b.desy.de; A.G. Akeroyd et al., hep-ex/0406071; I.I. Bigi and A.I. Sanda, hep-ph/0401003.

[2] S. Eidelman *et al.* [Particle Data Group Coll.], Phys. Lett. **B592**, 1 (2004) pdg.lbl.gov.

[3] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0408095; B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0406040; A. Garmash *et al.* [Belle Collaboration], Phys. Rev. D **69**, 012001 (2004) [arXiv:hep-ex/0307082]; S. Laplace [BABAR Collaboration], arXiv:hep-ex/0309049; K. Abe *et al.* [Belle Coll.], Phys. Rev. D **65**, 092005 (2002) [arXiv:hep-ex/0201007].

[4] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0407013; J. G. Smith, arXiv:hep-ex/0406063; B. Aubert *et al.* [BABAR Coll.], arXiv:hep-ex/0107075.

[5] S. Laplace and V. Shelkov, Eur.Phys.J.C 22, 431 (2001) (hep-ph/0105252).

[6] M. Suzuki, Phys. Rev. D 65, 097501 (2002).

[7] K. Maltman, Phys. Lett. **B462**, 14 (1999) [hep-ph/9906267]; Y. Meurice, Phys. Rev. D **36**, 2780 (1987); Mod. Phys. Lett. A **2**, 699 (1987); C. M. Shakin and H. Wang, Phys. Rev. D **63**, 074017 (2001); F. De Fazio and M. R. Pennington, Phys. Lett. B **521**, 15 (2001) [arXiv:hep-ph/0104289].

[8] see for example, S. Spanier and N.A. Tornqvist, Phys. Lett. B 592:1 (2004), C. Amsler and N.A. Tornqvist Phys. Rep. 389, 61 (2004) and references therein.

[9] F. E. Close and N. A. Tornqvist, J. Phys. G **28**, R249 (2002)[arXiv:hep-ph/0204205]; A. Abdel-Rehim, D. Black, A. H. Fariborz and J. Schechter, AIP Conf. Proc. **687**, 51 (2003); E. van Beveren and G. Rupp Mod. Phys. Lett. **A19**, 1949 (2004); M. Uehara hep-ph/0404221; M. R. Pennington, eConf **C030614**, 002 (2003) [arXiv:hep-ph/0310186]; J. L. Lucio Martinez and M. Napsuciale, Phys. Lett. B **454** (1999) 365 [arXiv:hep-ph/9903234]; J. L. Lucio Martinez and J. Pestieau, Phys. Rev. D **42** (1990) 3253; N. N. Achasov and V.N. Ivanchenko, Nucl. Phys. **B315** (1989)465; F.E. Close, N. Isgur, S. Kumano, Nucl. Phys. **B389**, 513 (1993); V. V. Anisovich and A. V. Sarantsev, Eur. Phys. J. A **16**, 229 (2003) [arXiv:hep-ph/0204328]; F. De Fazio and M. R. Pennington, Phys. Lett. B **521** (2001) 15 [arXiv:hep-ph/0104289].

[10] R. Akhmetshin *et al.* [CMD-2 Collaboration], Phys. Lett. B **462** (1999) 380 [arXiv:hep-ex/9907006]. V. M. Aulchenko *et al.* [SND Collaboration], Phys. Lett. B **440** (1998) 442 [arXiv:hep-ex/9807016]. M. N. Achasov *et al.*, Phys. Lett. B **485** (2000) 349 [arXiv:hep-ex/0005017]. J. E. Augustin *et al.* [DM2 Collaboration], Nucl. Phys. B **320** (1989) 1.

[11] E. M. Aitala *et al.* [E791 Collaboration], Phys. Rev. Lett. **89** (2002) 121801 [arXiv:hep-ex/0204018]; M. Ishida, S. Ishida, T. Komada and S. I. Matsumoto, Phys. Lett. B **518** (2001) 47; H. Muramatsu *et al.* [CLEO Collaboration], Phys. Rev. Lett. **89** (2002) 251802 [Erratum-ibid. **90** (2003) 059901] [arXiv:hep-ex/0207067].

[12] D. V. Bugg, Phys. Lett. B **572** (2003) 1 [Erratum-ibid. B **595** (2004) 556]; G. Colangelo, J. Gasser and H. Leutwyler, Nucl. Phys. B **603** (2001) 125 [arXiv:hep-ph/0103088]; R. Escribano, A. Gallegos, J. L. Lucio M, G. Moreno and J. Pestieau, Eur. Phys. J. C **28** (2003) 107 [arXiv:hep-ph/0204338], A. Gallegos, J. L. Lucio M. and J. Pestieau, Phys. Rev. D **69** (2004) 074033 [arXiv:hep-ph/0311133].

[13] J. A. Oller, E. Oset and J. R. Pelaez, Phys. Rev. D **59** (1999) 074001 [Erratum-ibid. D **60** (1999) 099906] [arXiv:hep-ph/9804209]; J. A. Oller, Nucl. Phys. A **727** (2003) 353 [arXiv:hep-ph/0306031].

[14] V. Chernyak, Phys. Lett. B **509**, 273 (2001) [arXiv:hep-ph/0102217].
A. C. Katoch and R. C. Verma, Int. J. Mod. Phys. A **11**, 129 (1996).

[15] see for examples, T. V. Brito, F. S. Navarra, M. Nielsen and M. E. Bracco, arXiv:hep-ph/0411233; A. Gokalp, Y. Sarac and O. Yilmaz, arXiv:hep-ph/0410380; H. G. Dosch, E. M. Ferreira, F. S. Navarra and M. Nielsen, Phys. Rev. D **65** (2002) 114002 [arXiv:hep-ph/0203225]. A. Ali, V. M. Braun and H. Simma, Z. Phys. C **63** (1994) 437 [arXiv:hep-ph/9401277]. V. M. Braun and A. V. Kolesnichenko, Phys. Lett. B **175** (1986) 485 [Sov. J. Nucl. Phys. **44** (1986 YAFIA,44,756-773.1986) 489.1986 YAFIA,44,756].

[16] P. Minkowski and W. Ochs, arXiv:hep-ph/0404194; arXiv:hep-ph/0304144.; Nucl. Phys. Proc. Suppl. **121**, 123 (2003) [arXiv:hep-ph/0209225]; Nucl. Phys. Proc. Suppl. **121**, 119 (2003) [arXiv:hep-ph/0209223]; Eur. Phys. J. C **9**, 283 (1999) [arXiv:hep-ph/9811518].
E. Klempt, B. C. Metsch, C. R. Munz and H. R. Petry, Phys. Lett. B **361**, 160 (1995) [arXiv:hep-ph/9507449].
E. Klempt, arXiv:hep-ex/0101031.
V. V. Anisovich and A. V. Sarantsev, Eur. Phys. J. A **16**, 229 (2003) [arXiv:hep-ph/0204328].

[17] C. H. Chen, Phys. Rev. D **67**, 094011 (2003) [arXiv:hep-ph/0302059]; Phys. Rev. D **67**, 014012 (2003) [arXiv:hep-ph/0210028]. H. Y. Cheng, Phys. Rev. D **67**, 034024 (2003)[arXiv:hep-ph/0212117].

[18] A. Ali, G. Kramer and C.D. Lu, Phys. Rev. D **58**, 094009 (1998) [arXiv:hep-ph/9804363]; A. Ali and C. Greub, Phys. Rev. D **57**, 2996 (1998) [arXiv:hep-ph/9707251].

[19] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Nucl. Phys. B **606**, 245 (2001) [arXiv:hep-ph/0104110]; G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996) [arXiv:hep-ph/9512380].

[20] Y.H. Chen, H.Y. Cheng, B. Tseng and K. C. Yang, Phys. Rev. D **60**, 094014 (1999) [arXiv:hep-ph/9903453].

[21] Y.Y. Keum and H. n. Li, Phys. Rev. D **63**, 074006 (2001) [arXiv:hep-ph/0006001]; Y.Y. Keum, H.N. Li and A.I. Sanda, Phys. Rev. D **63**, 054008 (2001) [arXiv:hep-ph/0004173]; Y.Y. Keum and A.I. Sanda, Phys. Rev. D **67**, 054009 (2003) [arXiv:hep-ph/0209014]; Y.Y. Keum, H. n. Li and A.I. Sanda, Phys. Lett. B **504**, 6 (2001) [arXiv:hep-ph/0004004].

[22] C.D. Lu, K. Ukai and M.Z. Yang, Phys. Rev. D **63**, 074009 (2001) [arXiv:hep-ph/0004213]; C.D. Lu and M.Z. Yang, Eur. Phys. J. C **23**, 275 (2002) [arXiv:hep-ph/0011238]; C.D. Lu and D.X. Zhang, Phys. Lett. B **400**, 188 (1997) [arXiv:hep-ph/9703439].

[23] M. Guagnelli, F. Palombi, R. Petronzio and N. Tantalo, Nucl. Phys. Proc. Suppl. **119**, 616 (2003) [arXiv:hep-lat/0209113]; D. Becirevic, arXiv:hep-ph/0310072.

[24] A. Abada, D. Becirevic, P. Boucaud, J. P. Leroy, V. Lubicz and F. Mescia, mesons from lattice QCD," Nucl. Phys. B **619**, 565 (2001) [arXiv:hep-lat/0011065]; D. Becirevic, S. Prelovsek and J. Zupan, Phys. Rev. D **68**, 074003 (2003) [arXiv:hep-lat/0305001]; D. Bećrević, S. Prelovšek and Ju. Zupan, Phys. Rev. **D67**, 054010 (2003); ICHEP-02, Amsterdam 2002; P. Ball and R. Zwicky, arXiv:hep-ph/0406261

[25] G. P. Lepage and S. J. Brodsky, Phys. Rev. D **22**, 2157 (1980); G. R. Farrar and D. R. Jackson, Phys. Rev. Lett. **43**, 246 (1979); S. Brodsky, Lett. Nuov. Cim. **7**, 719 (1973); S. J. Brodsky and G. R. Farrar, Phys. Rev. D **11**, 1309 (1975); Phys. Rev. Lett. **31**, 1153 (1973).

[26] D. s. Du, D. Yang and G. Zhu, Phys. Rev. D **60**, 054015 (1999).

[27] M. Diehl and G. Hiller, JHEP **0106**, 067 (2001) [arXiv:hep-ph/0105194].

[28] A. Deandrea and A. D. Polosa, Phys. Rev. Lett. **86**, 216 (2001) [arXiv:hep-ph/0008084].
N. Paver and Riazuddin, arXiv:hep-ph/0107330.

[29] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, arXiv:hep-ph/0407017.

[30] D. Black, A. H. Fariborz, F. Sannino and J. Schechter, Phys. Rev. D **59** (1999) 074026 [arXiv:hep-ph/9808415].

[31] M. Napsuciale and S. Rodriguez, Int. J. Mod. Phys. A **16** (2001) 3011 [arXiv:hep-ph/0204149].

ACKNOWLEDGEMENTS

The work of D.D. was supported by project PROMEP/103.5/04/1335 (México) and the work of J.L.M. was supported by CONACyT, México under project 37234-E. C.A.R. wants to thank the hospitality of J.L.M. and D.D. at IFUG, where part of this work was done.

APPENDIX

Below we list the $X^a_{b,c}$ expressed in terms of the form factors. We quote only those needed to compute the branching ratios given in the paper:

$$
\begin{aligned}
X^{\pi^-}_{\bar{B}^0 a_0^+} &= <\pi^-|(\bar{d}u)_L|0><a_0^+|(\bar{u}b)_L|\bar{B}^0>= f_\pi(m_B^2-m_a^2)F_0^{\bar{B}^0 a_0^+}(m_\pi^2) \\
X^{a_0^-}_{\bar{B}^0\pi^+} &= <a_0^-|(\bar{d}u)_L|0><\pi^+|(\bar{u}b)_L|\bar{B}^0>= -f_a(m_B^2-m_\pi^2)F_0^{\bar{B}^0\pi^+}(m_a^2) \\
X^{a_0^-}_{B^-\pi^0} &= <a_0^-|(\bar{d}u)_L|0><\pi^0|(\bar{u}b)_L|B^->= f_a(m_B^2-m_\pi^2)F_0^{B^-\pi^0}(m_a^2) \\
X^{\pi^0_u}_{B^-a_0^-} &= <\pi^0|(\bar{u}u)_L|0><a_0^-|(\bar{d}b)_L|B^->= \frac{f_\pi}{\sqrt{2}}(m_B^2-m_a^2)F_0^{B^-\pi^-}(m_\pi^2) \\
X^{\pi^-}_{B^-S^0} &= <\pi^-|(\bar{u}d)_L|0><S^0|(\bar{u}b)_L|B^->= f_\pi(m_B^2-m_{S^0}^2)F_0^{B^-S^0}(m_\pi^2) \\
X^{B^-}_{S^0\pi^-} &= <S^0P^-|(\bar{d}u)_L|0><0|(\bar{u}b)_L|B^->= -f_B(m_{S_0}^2-m_\pi^2)F_0^{S_0\pi^-}(m_B^2) \\
X^{B^-}_{a_0^-\pi^0} &= <a_0^-\pi^0|(\bar{d}u)_L|0><0|(\bar{u}b)_L|B^->= -f_B(m_a^2-m_\pi^2)F_0^{a_0^-\pi^0}(m_B^2) \\
X^{\bar{B}^0}_{\bar{K}^0S_0} &= <\bar{K}^0S^0|(\bar{s}d)_L|0><0|(\bar{d}b)_L|\bar{B}^0>= -f_B(m_{S^0}^2-m_K^2)F_0^{S^0\bar{K}^0}(m_B^2) \\
X^{B^-}_{S_0K^-} &= <S^0K^-|(\bar{s}u)_L|0><0|(\bar{u}b)_L|B^->= -f_B(m_S^2-m_K^2)F_0^{S^0K^-}(m_B^2) \\
X^{\bar{B}^0}_{(a_0^-\pi^+)_u} &= <a_0^-\pi^+|(\bar{u}u)_L|0><0|(\bar{d}b)_L|\bar{B}^0>= -f_B(m_{a_0^-}^2-m_\pi^2)F_0^{a_0^-\pi^+}(m_B^2) \\
X^{\bar{B}^0}_{a_0^+\pi^-} &= <a_0^+\pi^-|(\bar{u}u)_L|0><0|(\bar{d}b)_L|\bar{B}^0>= -f_B(m_{a_0^-}^2-m_\pi^2)F_0^{a_0^+\pi^-}(m_B^2) \\
X^{\bar{K}^0}_{\bar{B}^0S^0} &= <\bar{K}^0|(\bar{s}d)_L|0><S^0|(\bar{d}b)_L|\bar{B}^0>= f_{\bar{K}^0}(m_B^2-m_{S^0}^2)F_0^{\bar{B}^0S^0}(m_K^2) \\
X^{K^-}_{B^-S^0} &= <K^-|(\bar{u}s)_L|0><S^0|(\bar{u}b)_L|B^->= f_K(m_B^2-m_{S^0}^2)F_0^{B^-S^0}(m_\pi^2) \\
\tilde{X}^{a_0^-}_{\bar{B}^0\pi^+} &= <a_0^-|\bar{d}u|0><\pi^+|\bar{u}b|\bar{B}^0>= m_a\tilde{f}_{a_0^-}\frac{m_B^2-m_\pi^2}{m_b-m_u}F_0^{\bar{B}^0\pi^+}(m_a^2) \\
\tilde{X}^{S^0_d}_{B^-\pi^-} &= <S^0|\bar{d}d|0><\pi^-|\bar{d}b|B^->= \frac{m_{S_0}\tilde{f}_{S^0_d}}{m_b-m_d}(m_B^2-m_\pi^2)F_0^{B^-\pi^-}(m_{S_0}^2) \\
\tilde{X}^{a_0^-}_{B^-\pi^0} &= <a_0^-|\bar{d}u|0><\pi^0|\bar{u}b|B^->= m_a\tilde{f}_{a_0^-}\frac{m_B^2-m_\pi^2}{m_b-m_d}F_0^{B^-\pi^0}(m_a^2) \\
\tilde{X}^{S^0_s}_{B^-K^-} &= <S^0|\bar{s}s|0><K^-|\bar{s}b|B^->= m_{S^0}\tilde{f}_{S^0_s}\frac{m_B^2-m_K^2}{m_b-m_s}F_0^{B^-K^-}(m_{S^0}^2) \\
\tilde{X}^{S^0_s}_{\bar{B}^0\bar{K}^0} &= <S^0|\bar{s}s|0><\bar{K}^0|\bar{s}b|\bar{B}^0>= m_{S^0}\tilde{f}_{S^0_s}\frac{m_B^2-m_K^2}{m_b-m_s}F_0^{\bar{B}^0\bar{K}^0}(m_{S^0}^2) = \tilde{X}^{S^0_s}_{B^-K^-} \qquad (37)
\end{aligned}
$$

where S_0 is a neutral scalar (a_0^0 or f^0).

FORM FACTORS DEFINITIONS AND CONVENTIONS.

In order to compute the amplitude using the factorization, we use the following parametrization of the form factors. The decay constants are defined as:

$$\begin{aligned} \langle 0|A_\mu|P(q)\rangle &= i f_P q_\mu && (38)\\ \langle 0|\bar{q}_1\gamma_5 q_2|P(q)\rangle &\simeq \frac{-i f_P m_P^2}{m_1+m_2} \equiv \bar{f}_P m_P && (39)\\ \langle a_0^-|\bar{d}\gamma_\mu u|0\rangle &= f_{a0} p_\mu && (40)\\ \langle a_0^-|\bar{d}u|0\rangle &= m_{a0}\bar{f}_{a0} && (41) \end{aligned}$$

Using the equations of motion $(-i\partial^\mu(\bar{q}_1\gamma_\mu\gamma_5 q_2) = (m_1+m_2)\bar{q}_1\gamma_5 q_2$ and $-i\partial^\mu(\bar{q}_1\gamma_\mu q_2) = (m_1-m_2)\bar{q}_1 q_2$[18, 20] on can show that $\bar{f}_S = m_S f_S/(m_1-m_2)$ and that $f_{S^0}=0$ for a neutral scalar.

Form factors are defined as:

$$\begin{aligned} < M_2(p_2)|L_\mu|M_1(p_1) > &= \left(p_1+p_2-\frac{m_1^2-m_2^2}{q^2}q\right)_\mu F_+^{M_1M_2} + \frac{m_1^2-m_2^2}{q^2} q_\mu F_0^{M_1M_2}(q^2) && (42)\\ < M_2(p_2)M_1(p_1)|L_\mu|0 > &= \left(p_2-p_1-\frac{m_2^2-m_1^2}{q^2}q\right)_\mu F_+^{M_2M_1}(q^2) + \frac{m_2^2-m_1^2}{q^2} q_\mu F_0^{M_2M_1}(q^2) && (43) \end{aligned}$$

where $L_\mu = \gamma^\mu \frac{1-\gamma_5}{2} = \gamma^\mu P_L$. A factor of $-i$ has to be added to the form factors in the case one of the mesons is scalar.